

7

CHAPTER

BONDING IN ORGANIC MOLECULES

7.1 Petroleum Refining and the Hydrocarbons

7.2 The Alkane

7.3 The Alkenes and Alkynes


7.4 Aromatic Hydrocarbons

7.5 Fullerenes


7.6 Functional Groups and Organic Reactions

Connections to Biology: Functional Groups in Proteins

7.7 Pesticides and Pharmaceuticals



A petroleum refining tower
General Chemistry II



Cubane C₈H₈

7.1 PETROLEUM REFINING AND THE HYDROCARBONS


Petroleum: in latin ~ *petra (rock) + oleum (oil)* --- Crude Oil
Documented of usage 4000 yrs ago in Babylon.
Has been used as fuel in China since 400 B.C.
Has been used as a medicine since 15C. In Europe.
In 1854, world first modern oil well
In 1856, world first refinery
In 1859, world first actual modern oil well

Petroleum: major constituents are **hydrocarbons**

Hydrocarbon: compounds of hydrogen and carbon

Carbon - 83 to 87%, Hydrogen - 10 to 14%,

Nitrogen: 0.1 to 2%, Oxygen: 0.05 to 1.5%, Sulfur: 0.05 to 6.0%, Metals: < 0.1%




What can you make from one barrel of oil?

Researchers broke down a typical barrel of domestic crude oil into what could be produced from it. The average domestic crude oil has a gravity of **32 degrees** and weighs **7.21 pounds per gallon**. Here's what just one barrel of crude oil can produce:


Products Made from a Barrel of Crude Oil

Typical Products Made from a 42-Gallon Barrel of Refined Crude Oil




- 3% Asphalt
- 4% Liquefied Petroleum
- 10% Jet Fuel
- 18% Other Products
- 23% Diesel Fuel & Heating Oil
- 47% Gasoline

Source: U.S. Department of Energy.



There would be enough petrochemicals left in that same barrel to also provide the base for:




The lighter materials in a barrel are used mainly for paint thinners and dry-cleaning solvents, and they can make nearly a quart of one of these products. The miscellaneous fraction of what is left still contains enough byproducts to be used in medicinal oils, still gas, road oil and plant condensates.


It's a real industrial horn of plenty.

7.2 THE ALKANES


- **Hydrocarbons** – compounds that are only composed of **hydrogen** and **carbon**




Ethane
 C_2H_6



Ethylene
 C_2H_4



Acetylene
 C_2H_2



Benzene
 C_6H_6



- Which of the molecules above is saturated with hydrogen atoms?

7.2 THE ALKANES



◆ Normal Alkanes: straight chain alkanes

Alkanes: Saturated Hydrocarbons

CH_4 : sp^3 hybridized



C_2H_6




Saturated hydrocarbon : *all bonds are single bonds*




“Alkane” paraffin

| | |
|------------------------------|-------------|
| CH_4 | methane |
| C_2H_6 | ethane |
| C_3H_8 | propane |
| C_4H_{10} | butane |
| C_5H_{12} | pentane |
| C_6H_{14} | hexane |
| C_7H_{16} | heptane |
| C_8H_{18} | octane |
| $\text{C}_{30}\text{H}_{62}$ | triacontane |


b.p. & m.p.
increase




◆ Wax



Polyethylene



Polymerization



Ethylene
(monomer)

Polyethylene
(polymer)



Saturated hydrocarbon : *all bonds are single bonds*

“Alkane” *paraffin*




of structures --- *isomers*


Branched-Chain Alkanes & Isomerism

| | | | |
|---------------------------------|-------------|------------------------|--|
| CH ₄ | methane | 1 | |
| C ₂ H ₆ | ethane | 1 | |
| C ₃ H ₈ | propane | 1 | |
| C ₄ H ₁₀ | butane | 2 | |
| C ₅ H ₁₂ | pentane | 3 | |
| C ₆ H ₁₄ | hexane | 5 | |
| C ₇ H ₁₆ | heptane | 9 | → beginning of stereoisomerism (11) |
| C ₈ H ₁₈ | octane | 18 | |
| C ₃₀ H ₆₂ | triacontane | 4.11 x 10 ⁹ | |


Example : C_5H_{12}



Pentane : b.p. $36^{\circ}C$












Isopentane : b.p. $28^{\circ}C$











Neopentane : b.p. $9.5^{\circ}C$

example : C_7H_{16} : Heptane



9



◆ Isomerism:


1. Structural (constitutional) isomer: different bonding arrangements of the same atoms.

구조 이성질체

2. Stereoisomer: same bonding arrangement, different spatial positions.

입체 이성질체



-- enantiomer (거울상 이성질체)




These are mirror images and not superimposable
i.e. different compounds.

Different but have same physical properties except optical rotation.
i.e. inseparable

Stereoisomers



enantiomers



enantiomers

◆ Cyclic Alkanes:


Saturated hydrocarbon : *Ring structures*




Cyclic Compounds

*Usually unstable when it's small
Associate with **strain energy***


Most stable cyclic compound



Cyclohexane



chair form



boat form



Cyclopropane

*Most strained
reactive*



Cyclopropane

- Cyclopropane is 44 kJ/mol less stable than cyclohexane per CH_2 group. It is highly strained and very reactive

1. Angle strain


- Bond angles of 60° cause electron pair repulsion in adjacent bonds
- Inefficient sigma bond overlap

2. **Torsional strain** – eclipsing C-H bonds all the way around the ring








7.3 THE ALKENES AND ALKYNES

Alkenes



ALKYNES



The Alkenes and Alkynes

➤ Unsaturated hydrocarbons

Alkene ~ double bonds ex. Ethene (Ethylene), C_2H_4

Alkyne ~ triple bonds ex. Ethyne (Acetylene), C_2H_2







Fig. 7.10. Reaction with $KMnO_4$.
(a) No reaction with hexane.
(b) Redox reaction with 1-hexene.
Products: MnO_2 and
 $CH_3(CH_2)_3CH(OH)CH_2OH$

Unsaturated hydrocarbon : *Alkenes, Alkynes*


Alkene : C_nH_{2n}



Pi bonds are more reactive than sigma bonds




Trans isomer is more stable due to steric effect




2. Stereoisomer: same bonding arrangement, different spatial positions.

입체 이성질체

geometrical isomer



trans




cis

rotation around the double bond requires **60 kcal/mol**

Different compounds different physical, chemical properties







p- bonding






High-energy structure

Unsaturated hydrocarbon : *Alkenes, Alkynes*




Unsaturated hydrocarbon : *Alkenes, Alkynes*



Polyenes





p molecular orbitals of butadiene; see chapter 20 p976



Conjugated p bonding



153 pm




*sp*³ *sp*³

More stable than isolated two double bonds

7.4 AROMATIC HYDROCARBONS



Hydrocarbons with $C_{4n+2}H_{2n+4}$

Benzene: simplest example C_6H_6



Modern view of three double bonds: *delocalized*

p molecular orbitals; view from the top




represented as circle inside


more stable than trienes
i.e. much less reactive

Copyright © 2006 Pearson Education, Inc., Publishing as Benjamin Cummings


Aromatics from petroleum




naphthalene




toluene







***o*-xylene**



***m*-xylene**



***p*-xylene**



7.6 FUNCTIONAL GROUPS AND ORGANIC REACTIONS


TABLE 18.4

Common hydrocarbon derivatives


| Derivative | Functional group | General formula | Examples | |
|-----------------|---|---|--|--|
| Halide | $-\text{Cl}, -\text{Br}$ | $\text{R}-\text{Cl}$ | CH_2Cl_2 , methylene chloride (dichloromethane) | $\text{CH}_3\text{CHClCH}_3$ isopropyl chloride (2-chloropropane) |
| Alcohol | $-\text{OH}$ | $\text{R}-\text{OH}$ | CH_3OH methanol | $\text{CH}_3\text{CH}_2\text{OH}$ ethanol |
| Ether | $-\text{O}-$ | $\text{R}-\text{O}-\text{R}'$ | $\text{CH}_3\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_3$ Diethyl ether | $\text{CH}_3-\text{O}-\text{C}(\text{CH}_3)_3$ Methyl <i>t</i> -butyl ether (MTBE) |
| Ketone | $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}- \end{array}$ | $\begin{array}{c} \text{O} \\ \parallel \\ \text{R}-\text{C}-\text{R}' \end{array}$ | $\begin{array}{c} \text{O} \\ \parallel \\ \text{CH}_3-\text{C}-\text{CH}_3 \end{array}$ Acetone (propanone) | $\begin{array}{c} \text{O} \\ \parallel \\ \text{CH}_3-\text{C}-\text{CH}_2\text{CH}_3 \end{array}$ Methyl ethyl ketone (MEK) (butanone) |
| Aldehyde | $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}-\text{H} \end{array}$ | $\begin{array}{c} \text{O} \\ \parallel \\ \text{R}-\text{C}-\text{H} \end{array}$ | $\begin{array}{c} \text{O} \\ \parallel \\ \text{H}-\text{C}-\text{H} \end{array}$ Formaldehyde (methanal) | $\begin{array}{c} \text{O} \\ \parallel \\ \text{CH}_3-\text{C}-\text{H} \end{array}$ Acetaldehyde (ethanal) |
| Carboxylic acid | $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}-\text{OH} \end{array}$ | $\begin{array}{c} \text{O} \\ \parallel \\ \text{R}-\text{C}-\text{OH} \end{array}$ | $\begin{array}{c} \text{O} \\ \parallel \\ \text{CH}_3-\text{C}-\text{OH} \end{array}$ Acetic acid (ethanoic acid) | $\begin{array}{c} \text{O} \\ \parallel \\ \text{CH}_3\text{CH}_2-\text{C}-\text{OH} \end{array}$ Propionic acid (propanoic acid) |




Alkyl halides



Methylene
chloride



Chloroform




Carbon
tetrachloride




1. Alcohol : $\text{R}-\text{OH}$

2. Ether : $\text{R}-\text{O}-\text{R}'$



Diethyl ether -- "Ether"




MTBE

KAIST
CHEMISTRY




3. Aldehyde, Ketone :




R' = H : aldehyde

4. Carboxylic Acid :




5. Ester :




Alcohol


R-OH




Methanol




Ethanol
(primary alcohol)




iso-Propanol
(secondary alcohol)




Ethylene Glycol




Glycerol




t-butyl alcohol



Menthol




Phenol




Urushiol


Aldehyde, Ketone




Formaldehyde



Acetone




Acetaldehyde




Benzaldehyde
(cherry flavor)


Carboxylic acid




Formic acid




Acetic acid



Benzoic acid




Ibuprofen (부루펜)




Naproxen (낙센)


Ester




Pear




Banana




Pineapple




Apple



Jasmine






- Product of the reaction between a carboxylic acid and an alcohol
- Fragrant odors, flavors of fruits



Tristearin, $\text{C}_{57}\text{H}_{110}\text{O}_6$: animal fat

Triglycerides


| STRUCTURE AND NAME | NUMBER OF CARBON ATOMS | NUMBER OF CARBON-CARBON DOUBLE BONDS | MELTING POINT (°C) |
|--|------------------------|--------------------------------------|--------------------|
| SATURATED  Myristic acid | 14 | 0 | 54 |
|  Stearic acid | 18 | 0 | 69 |
| UNSATURATED  Linoleic acid | 18 | 2 | -5 |

Trans fats (트랜스지방脂肪)


- ❖ Hydrogenation of oils (ester of cis-unsaturated fatty acids)
 - Saturated fats with higher m. p.:
 - solid, good for baking and extended shelf-life
 - Remaining double bonds converted from cis to trans isomers → bad for health!



●Amine●




Cadaverine
(시체썩는 냄새)




Methamphetamine
(필로폰)


●Amide●



Nylon-6





Acetaminophen
(타이레놀)




Lidocaine
(국소마취제)



Stereoisomers




enantiomers




enantiomers

Stereoisomers




Two chiral centers make the situation complicated !



Diastereomers: Stereoisomers are not mirror images of one another and non-superimposable.


Stereochemistry




- Separation of enantiomers
- Recognition of enantiomers differently

Importance of Stereochemistry

Limonene




lemon




orange

Aspartame




sweet




bitter

Thalidomide




sedative



teratogenic


Reactions of organic compounds

1. carbon-carbon bond formation & cleavage



2. general reactive intermediates

General Chemistry II








KAIST
CHEMISTRY










Conversion of organic functional groups

● Oxidation




● Reduction




Synthesis & reactions of Esters

How does esters form?

Fischer Esterification



Synthesis of Amide is similar




7.7 PESTICIDES AND PHARMACEUTICALS

Impact of Organic Compounds to the World


➤ 90% of matter on earth are organic!

i.e. organic compounds are everywhere around us.


In food




Sucrose (sugar)




MSG (glutamic acid)




Vitamin C




Citric acid (Lemon)




beta-carotene (carrot)




Cyanidin (anthocyanin)




Tartaric acid (grape)



Capsaicin (pepper)




Isothiocyanate (wasabi)




Cinnamaldehyde (cinnamon)

General Chemistry II


From nature (natural products) and beyond.....




Pyrethrin I (from chrysanthemum)
weak insecticide




Dioxin




Morphine




Cocaine




Permethrin (synthetic pyrethrin)




TCDD
(고엽제성분)




penicillin




atropine




nicotine




imidacloprid




VX




estrogen




cholesterol




DDT




malathion



Glyphosate (weed killer)




Dianabol




testosterone

More natural products




R = 5'-deoxyadenosyl, Me, OH, CN


Vitamin B12




quinine




artemisinin




Tetrodotoxin (fugu fish, 복어)




**Avermectin
(심장사상충)**



Ginkgolide (은행)




brevetoxin (red tide, 적조)




Maitotoxin, C₁₆₄H₂₅₆O₆₈S₂Na₂, M.W. = 3422


Significant organic molecules made by chemists




Mauveine (dye, purple)




Nylon 66




polystyrene




teflon




OLED materials




Prontosil (antibiotic)




Norethisterone (contraceptive)




viagra



Omeprazole (antiulcer)



Olanzapine (schizophrenia)



Prozac (antidepressant)